

DRUG FACTS

Active ingredient(s)

Purpose

Use(s)

Warnings

Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers

Do not use

- with other acid reducers
- if you have kidney disease, except under the advice and supervision of a doctor
- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- nausea or vomiting
- stomach pain
- unexplained weight loss
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Ask a doctor or pharmacist before use if you are

When using this product

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding

Keep out of reach of children

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Other information

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by certain foods and beverages
- adults and children 12 years and over:
- to **relieve** symptoms, swallow 1 tablet with a glass of water
- to **prevent** symptoms, swallow 1 tablet with a glass of water **30 to 60 minutes before** eating food or drinking beverages that cause heartburn
- can be used up to twice daily (do not take more than 2 tablets in 24 hours)
- do not chew tablet
- children under 12 years: ask a doctor
- Blister: do not use if individual blister unit is open or torn
- Bottle: do not use if printed foil under bottle cap is open or torn
- store at 20°-25°C (68°-77°F)
- avoid excessive heat or humidity
- this product is sodium and sugar free

Dist: Boehringer Ingelheim Consumer Health Care Products Division of Boehringer Ingelheim Pharmaceuticals, Inc. Ridgefield, CT 06877 ©2009

Inactive ingredients

carrageenan, FD&C blue no.1, flavors, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol,polysorbate, sucralose, and titanium dioxide